



Citation for published version:

Trewartha, G, Preatoni, E, England, ME & Stokes, KA 2015, 'Injury and biomechanical perspectives on the rugby scrum: a review of the literature', *British Journal of Sports Medicine*, vol. 49, no. 7, pp. 425-433.
<https://doi.org/10.1136/bjsports-2013-092972>

DOI:

[10.1136/bjsports-2013-092972](https://doi.org/10.1136/bjsports-2013-092972)

Publication date:

2015

Document Version

Peer reviewed version

[Link to publication](#)

University of Bath

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **TITLE PAGE**

2 **Title**

3 Injury and biomechanical perspectives on the rugby scrum: a review of the literature

4 **Author list**

5 Grant Trewartha^(a,✉), Ezio Preatoni^(a), Michael E. England^(a,b) and Keith A. Stokes^(a)

6 **Affiliations**

7 ^(a) Sport, Health and Exercise Science, Department for Health, University of Bath, UK

8 ^(b) Rugby Football Union, Twickenham, UK

9 ✉ **Corresponding author**

10 Grant Trewartha

11 g.trewartha@bath.ac.uk

12 +44 (0)1225 383055

13 Sport, Health & Exercise Science – Department for Health, University of Bath

14 Applied Biomechanics Suite, 1.301

15 BA2 7AY - BATH (UK)

16 **Keywords**

17 Injury epidemiology, biomechanics, injury mechanisms

18 **Word count:**

19 7861

20

21 **Contributorship Statement**

22 Grant Trewartha: is guarantor. Initiated the overall project and supervised all its phases;
23 performed the primary literature searches; drafted the paper; contributed to revising the
24 paper; approved the final version of the paper.

25 Ezio Preatoni: contributed to the provision of grey literature; drafted the 'Mechanisms of
26 injury' section; contributed to revising the paper; approved the final version of the paper.

27 Mike England: contributed to the provision of grey literature; contributed to revising the
28 paper; approved the final version of the paper.

29 Keith Stokes: contributed to the provision of grey literature; contributed to revising the paper;
30 approved the final version of the paper.

31

32 **What this study adds**

- 33 - Synthesises the recent research literature pertaining to the medical and
34 biomechanical aspects of rugby union scrummaging
- 35 - Highlights the need to consider both acute catastrophic and chronic degenerative
36 injury types when considering injuries occurring in the rugby scrum
- 37 - Highlights that the most contemporary literature available is collectively confirming
38 that the forces involved in rugby scrummaging are high and multi-planar, but can be
39 modified appropriately by alterations to the scrum engagement technique

40

ABSTRACT

As a collision sport, Rugby Union has a relatively high overall injury incidence, with most injuries being associated with contact events. Historically, the set scrum has been a focus of the sports medicine community due to the perceived risk of catastrophic spinal injury during scrummaging. The contemporary rugby union scrum is a highly dynamic activity but to this point has not been well characterised mechanically. In this review we synthesise the available research literature relating to the medical and biomechanical aspects of the rugby union scrum, in order to 1) review the injury epidemiology of rugby scrummaging; 2) consider the evidence for specific injury mechanisms existing to cause serious scrum injuries; and 3) synthesise the information available on the biomechanics of scrummaging, primarily with respect to force production. The review highlights that the incidence of acute injury associated with scrummaging is moderate but the risk per event is high. The review also suggests an emerging acknowledgement of the potential for scrummaging to lead to premature chronic degeneration injuries of the cervical spine and summarises the mechanisms by which these chronic injuries are thought to occur. More recent biomechanical studies of rugby scrummaging confirms that scrum engagement forces are high and multi-planar, but can be altered through modifications to the scrum engagement process which control the engagement velocity. Because the set scrum is a relatively “controlled” contact situation within Rugby Union it remains an important area for intervention with a long term goal of injury reduction.

INTRODUCTION

Rugby union (rugby) is a contact sport involving periods of submaximal activity such as walking and jogging interspersed with short bouts of high intensity activity such as sprinting, and game-specific events such as the tackle, maul, ruck and scrum.¹ Rugby union has comparatively high injury incidence,²⁻⁵ albeit similar to other collision sports.⁶⁻⁹ There has been a focus on the safety of specific contact elements of the game, with the scrum and the tackle at the forefront as the game events the International Rugby Board has targeted for injury prevention initiatives. The set scrum is of particular interest, since as a set piece phase there is a view that injury occurrence from the scrum should be to some extent “controllable”.¹⁰ Nevertheless, the scrum is perceived as a phase of play with considerable injury risk, particularly in the context of the risk of chronic or catastrophic spinal injury. This is principally because, although catastrophic injuries due to scrummaging are very rare,¹¹ they are exceptionally debilitating. Although some commentators have called for a radical alteration of the scrum or even its ban from certain levels of rugby to reduce this perceived injury risk,¹² these views have been equally strongly rebuffed.¹¹

According to Law 20 of the International Rugby Board (IRB) Law Book which governs the rugby scrum, the purpose of the scrum is to “restart play quickly, safely and fairly, after a minor infringement or a stoppage”.¹³ A scrum involves a maximum of 8 players per team (the forward pack), who bind together in three rows (front, second and back), and then bind with an opposition forward pack to compete for possession of the ball by exerting a coordinated pushing action (Figure 1). Effective scrummaging requires a pack of forwards to produce forceful and coordinated actions to ensure dominance over the opposition, to provide a platform for launching attacks, and to disrupt opposition ball. Mechanically, contemporary scrummaging is broadly characterised by a high initial impact during the engagement of the opposing packs which is followed by the application of sustained opposing forces.^{14,15}

**** Figure 1 here ****

The rugby scrum generates very high biomechanical demands on players' musculoskeletal structures and thus exposes forwards, and front row forwards in particular, to the risk of both acute and chronic (overuse) injuries. The epidemiological studies of rugby injury reviewed later describe a moderate incidence/proportion of scrum-related injuries, but also the potential seriousness of these occurrences. In fact, even though some recent data suggest a relative decline in scrum-related serious injuries over recent decades, about 40% of all catastrophic (typically spinal cord) injuries that occur in rugby are related to scrummaging.¹⁶ Furthermore, players may appear asymptomatic in the shorter-term, but may experience repeated micro-trauma¹⁷ that contribute to the emergence of long-term degeneration and pathologies of the spine, including physical abnormalities^{18,19}, reduced mobility²⁰, and impaired proprioception.²¹

While rugby scrums may be associated with a number of potential injury risk factors, there is currently very little quantitative data to identify and describe these risk factors. There is a lack of information about the forces and motions involved in live contested scrummaging, and, consequently, little objective knowledge about how performance could be optimised and injuries prevented. Quantitative research on the rugby scrum has been occasional^{14,22-24} and has demonstrated that high forces can be generated, particularly during the engagement phase.¹⁴ Recent research has increased the scale and scope of the biomechanical investigation of the scrum,^{15,25} and has moved the investigation into live contested scrummaging to improve the ecological validity of the data.²⁶

The aim of this review is to synthesise the available literature relating to injury and biomechanical aspects of the rugby union scrum, and to 1) review the injury epidemiology of rugby scrummaging; 2) consider the evidence for specific injury mechanisms existing to cause serious scrum injuries; and 3) synthesise the information available on the biomechanics of scrummaging, primarily with respect to force production.

115

116 **METHODS**

117 **Search Strategy**

118 A literature search was conducted through Web of Knowledge and PubMed databases in
119 June 2011 and in March 2013. The search parameters were for years 1960-2011 and 2010-
120 2013 for the initial search and follow-up search, respectively. The search expressions were
121 'rugby and scrum', 'scrum and injury', 'rugby and spine', 'rugby and biomechanics', 'scrum
122 and biomechanics', 'rugby and injury prevention', and 'rugby and injury mechanism'. The
123 literature search was restricted to English and Italian language publications. The search
124 returned a total of 997 publications via Web of Knowledge and 1617 publications via
125 PubMed. The records returned were browsed for relevance via the title and abstracts, with
126 duplicate records being removed. The reference lists of included key studies, and relevant
127 "grey literature" (e.g. conferences proceedings) were manually searched to identify
128 additional articles.

129

130 **Selection Criteria**

131 Studies were selected for further review based on focus of study and population studied,
132 with case reports being given low priority. 137 studies were initially fully critiqued and
133 considered for inclusion in the manuscript.

134

135 **RESULTS AND DISCUSSION**

136 **Epidemiology of all-outcome scrummaging injuries**

137 The epidemiology of general rugby injury is well documented elsewhere.²⁷ Overall, elite
138 rugby union has comparatively high injury incidence²⁻⁵ in relation to other team sports, which
139 decreases as the playing level moves from the elite level to the community game²⁸⁻³⁰ and

youth levels.³¹⁻³⁴ Injury incidence also appears to be lower in women's compared with men's rugby.^{35,36}

Approximately 6-8% of all rugby injuries result from scrummaging, which is moderate compared with other injury-causing match events such as tackles.^{2-4,37} Brooks et al² found that scrummaging accounted for 11% of injuries to forwards, and the incidence of scrummaging injuries referenced to hours of exposure (10/1000 player match hours in elite senior rugby^{2,38}; ~2/1000 player match hours in youth rugby³⁹) is correspondingly moderate to low. However, when expressed as injury per event (propensity), scrum injuries are higher than for any other contact event, reported at 8.1 injuries/1000 scrums in English Premiership rugby.⁴⁰ Taking severity into account, the scrum has the highest injury risk per event of any contact event, with 213 days lost per 1000 scrum events. This is nearly double the risk per event of injuries from legal tackles. The risk of injury due to collapsed scrums versus completed scrums has also been found to be significantly higher at professional ($P=0.04$)³⁸ and community ($P<0.001$) level.⁴¹

It is important to note that a wide range of injuries are associated with scrummaging. In a cohort of professional rugby union players, calf muscle injuries were the most common scrummaging injury followed by lumbar spine injury, with calf muscle injuries and shoulder injuries causing the greatest number of days absence due to scrummaging.⁴² Neck injuries made up only 15% of the scrummaging injury burden. Front row forwards sustained 91% of all scrummaging injuries. Furthermore, the scrum was responsible for a high proportion of front row spinal injuries (41% of cervical; 56% of thoracic; 71% of lumbar).⁴² The reason for front row players' susceptibility to spinal injuries has been suggested to be the repeated high forces experienced by these players,²³ particularly during scrum engagement.

In comparing the injury profiles of each forward position versus all other forward playing positions, Brooks & Kemp⁴³ found that player absence due to neck injuries for loose head props and hookers was higher than other forward positions due mainly to cervical disc / nerve root injuries. These injuries were sustained mainly during tackling (57% for loose

head, 38% for hooker) but also scrummaging (29% for loose head, 19% for hooker). Possibly based on specificity of positional roles, the pattern of injury differed across the front row. Loose head props had more absence than other forwards due to shoulder rotator cuff injuries, primarily suffered to the right shoulder during scrummaging (66% of rotator cuff injuries). Tight head props had a greater absence due to lumbar spine injuries (67% of lumbar disc / nerve root injuries and 57% of lumbar soft tissue were attributed to scrummaging) and also due to calf injuries which were suffered mainly (54%) during scrummaging.

In summary, scrummaging accounts for a moderate proportion of the overall injury burden within rugby union, but scrummaging can be considered a high risk event in comparison with other game activities, particularly if the scrum collapses. Front row players are particularly susceptible to scrummaging injury and the scrum is responsible for a considerable proportion of the spinal and shoulder injuries sustained by front row forwards. In the context of the scrum being a relatively “controllable” event when compared with other match events such as the tackle, it is reasonable to suggest that there should be further efforts to reduce the injury burden of scrummaging.

Catastrophic Spinal Injuries in Rugby

Magnitude

In rare circumstances, a sports injury can result in permanent paralysis or a fatality. Spinal cord injuries resulting in fatal or catastrophic consequences cause significant concern in collision sports, such as American Football ⁴⁴ and Rugby Union. ¹⁶ There are difficulties in the definition of “serious spinal cord injuries” or “catastrophic” injuries, and studies do not always provide a definition of which injuries are covered. For the purposes of this review, serious and catastrophic will be used synonymously and relate to an injury resulting in neurological impairment without a return to full function, equivalent to ASIA classification of A

to D and therefore in line with the operational definition of catastrophic injury employed by the International Rugby Board.⁴⁵

Estimates suggest that the incidence of catastrophic spinal injury from rugby union may lie anywhere from 1-2/100,000 players per year to 10/100,000 players per year.^{16,46-50} This reflects a small number of injuries in the context of the playing population, but preventative strategies must be prioritised towards injuries causing permanent disability or death due to the devastating consequences of such injuries.⁵¹

Fuller¹¹ performed a risk analysis for sustaining a catastrophic spinal cord injury in rugby union and compared this with other collision sports and other common activities. The overall risk of catastrophic injury from rugby union ranged from 'acceptable' to 'tolerable' (as defined by the UK's Health and Safety Executive) depending on the country analysed. The risks in rugby union were described as *similar* or *less than* other sports such as American Football, rugby league and ice hockey, comparable to work-based risks and less than risks for motorcyclists and pedestrians. While acknowledging the subjective nature of perceiving risk in terms of the activity context, this study concluded that the risk of sustaining a catastrophic injury from rugby union was acceptable and the laws of the game adequately managed this risk, although all reasonably practicable measures should be taken to further reduce the risk in accordance with accepted risk management principles.

Match Event

Understanding which phase of play is considered responsible for causing catastrophic spinal injuries is a key step in prevention. For the period 1956-2004, the scrum was implicated as the match event causing a catastrophic injury in 42% of cases, compared with the tackle (34%), rucks/mauls (20%) and other phases (4%).¹¹ A separate review of published data for the period 1970-2001 offered similar findings, with scrums associated with approximately 40% of all serious cervical spine injuries in rugby union and the tackle associated with 36% of injuries.¹⁶ Bohu et al⁵² stated that 19 out of 37 recorded acute spinal cord injuries (i.e.,

51%) occurred in the scrum, although it wasn't obvious which other game events were associated with the remaining injuries.

There is some evidence to suggest that the tackle phase is becoming the game event most implicated in serious spinal cord injury.¹⁶ In South Africa during the period 1980-2007, 45% of 126 serious acute spinal cord injuries were attributed to the tackle phase, and 37% to the scrum.⁵³ In Australian rugby during the 1997-2002 period, 9 of 23 injuries occurred as a result of the tackle, 7 as a result of the scrum and 6 the ruck/maul.⁵⁴

A recent study by Brown and colleagues⁴⁶ updated the rugby-related catastrophic injury landscape in South Africa. In the period 2008-2011 45 acute spinal cord injuries were recorded, including near-miss events, resulting in an estimated annual incidence of 1.73 injuries per 100,000 players. The scrum accounted for 42% of these injuries (19 of 45) and the tackle 38% of injuries. There was a greater preponderance for scrum injury to occur to senior players and for the injuries caused by scrummaging to result in permanent disability. Information has recently been collected regarding admissions of under 19 rugby players to spinal units in Great Britain and Ireland between 1996 and 2010.⁵⁵ Thirty six injuries were recorded, 13 of which were associated with injuries in the scrum, compared with 17 associated with the tackle. The proportion of cases with complete neurological deficit following an injury in the scrum (61%) was significantly greater than in injuries following the tackle (29%, $P<0.001$). Overall, these findings suggest that acute spinal cord injuries occur at a similar frequency in the scrum and tackle, but that neck injuries sustained in the scrum are more likely to be more serious.

Trend over Time

It is difficult to ascertain whether the incidence of rugby-related serious cervical spine injuries has changed over the last 30 years due to a lack of accurate exposure (player numbers) data,¹⁶ because the raw number of injuries is relatively low, and because there are substantial differences in data collection methodologies. In Australia, there have been a

number of studies which overall suggest a tendency for a slight decrease in the incidence of serious spinal injuries over time. Taylor et al reported that the incidence of serious spinal cord injuries dropped from 4.6/100,000 players (1983-1989) to 3.0/100,000 players (1990-1996),⁵⁰ and was at 3/100,000 players for all football codes in 1997-2002.⁵⁴ Between 1975-1985 to 1986-1996 there was a 67% reduction in serious spinal cord injuries attributed to scrum engagement, from 12 injuries to 4 injuries,⁵⁰ and there was a tendency for acute spinal cord injuries to be less severe in the period 1997-2002 compared with 1986-1996.⁵⁴ Similar findings are reported by Berry et al⁵⁶ who tracked the incidence rates of severe cervical spinal cord injury in rugby union and rugby league over 17 years (1986-2003) found a non-significant decrease in the incidence rate of these severe injuries over time but with wide confidence intervals.

In French rugby, Bohu et al⁵² suggested a reduction in serious cervical spine injury incidence when comparing 1995-2001 (2.1/100,000 players) with 2001-2006 (1.4/100,000 players), and primarily attributed this to reduced incidence of injuries from scrummaging. The authors considered these reductions to be due to a change in scrum laws (e.g. limited engagement and pushing distances permitted at lower levels) and use of a 'front row forward passport' as medical clearance to play in these positions.

In South Africa, there was an apparent 48% reduction in the incidence of serious spinal injuries in schoolboys in the years 1990-1997 compared with previous datasets.⁵⁷ However, there was a 22% increase in admissions to spinal units in adult players in the 1990-1997 period compared with the 1982-1989 period,⁵⁷ echoing the results of Scher.⁵⁸ A retrospective pooled analysis, showed that the incidence of serious spinal cord injuries in South African rugby (assuming relatively consistent player numbers) between 1980-2007 had neither increased nor decreased, consistently lying somewhere between 0.5-1.0/100,000 players per year.⁵³

Overall, it is unclear whether there have been any changes over time in the rate of rugby-related catastrophic injuries, perhaps due to cultural, medical resource and reporting

differences and the fact that the absolute numbers of injuries are low. However, pooling data suggests there may have been slight reductions over the last 30 years.

Effect of Age and Playing Experience

Reports on the relative risk of a serious spinal injury in young and adult players appear contradictory,¹⁶ with some studies reporting younger players to be at higher risk^{59,60} and others suggesting adult players are at relatively higher risk.^{46,50,57,58,61} Noakes et al conducted a retrospective analysis on schoolboy and adult players in South African rugby and found that 80% of the 67 recorded serious spinal injuries occurred to adults and 20% to schoolboys.⁵⁷ No player numbers were reported but it was considered likely that there were more schoolboy than adult players. These findings are supported by Scher⁵⁸ who estimated that adult players were at 10-12 times greater risk of a serious spinal injury than schoolboy players, and Taylor et al⁵⁰ who showed serious SCI incidence of 6.9/100,000 players in adult players and 1.2/100,000 players in schoolboy rugby. Recent data from South Africa which has used estimates of playing populations has confirmed that adult players (5.3/100,000 players) are at increased risk of acute spinal cord injuries compared with junior players (0.9/100,000 players), with the injuries to adult players also being more likely to result in permanent impairments.⁴⁶

In older adult players, degenerative arthritis of the spine may be an additional risk factor for acute injury,^{17,19,62} although it can be argued that a lack of maturity in skeletal and ligamentous structures is a potential additional risk factor for younger players.⁶³ In terms of physical conditioning characteristics, there is no compelling evidence to suggest that body anthropometrics or training status is a major risk factor for spinal injury but most recommendations continue to advocate the need for suitable physical build and specific training for those players involved in scrummaging, particularly in the front row.

A mismatch in skill, experience or strength has been suggested as a risk factor for injury in the scrum, with the risk of injury being equal across the stronger and weaker team. Wetzler

et al ⁶⁴ found evidence of a mismatch of some type in 25% of all serious scrum injuries. These sentiments are echoes of other research or opinion pieces (e.g. ^{65,66}). Also, a lack of experience of playing in the front row has been highlighted previously as a risk factor for injury (e.g attributed in 39% of scrum injuries ⁵⁰) although this practice should now be impossible if IRB laws are enforced which state: “Each player in the front row and any potential replacement(s) must be suitably trained and experienced”. What constitutes the minimum standard for suitable training and experience and how this is monitored is likely to vary between different national unions and playing levels.

Playing Position

There is consistent evidence to show that front row forwards, and particularly hookers, are at highest risk for serious spinal cord injuries (Table 1). Hookers represent 7% of the players in a team, yet in South Africa, ⁴⁶ hookers account for 46% (12 of 26 injuries) of all the permanent outcome acute spinal cord injuries, with 83% (10 injuries) of these injuries occurring in the scrum. The vulnerability of the hooker in the scrum has been attributed to a number of factors, including the wrapping of their arms around props in the scrum with the effect that he or she cannot control or dissipate forces of engagement, the reliance on the props for support during engagement and formation, and the inability to adjust upper body position to react to improper engagement.

320

321 Table 1. Playing positions sustaining acute spinal cord injuries.

Study	Number of injuries	Percentage of injuries sustained by playing groups (%)		Percentage of injuries sustained by specific playing positions (%)			
		Forwards	Backs	Prop	Hooker	Other Forwards	Backs
Silver ⁶³	19	79	21				
Hermanus ⁵³	139	76	24	16	30	30	24
Bohu ⁵²	37	89	11	19	38		
Quarrie ^{16 **}	341	76	24	18	33	25	24
Brown ⁴⁶	40	86	14	13	38	35	14
Brown ^{46 *}	26	100	0	12	46	42	0

322 * Permanently disabling injuries included only

323 ** Only including injuries in the sample with known playing positions

324

325

Non-catastrophic Traumatic Spinal Injuries

Most injuries to the spine from rugby are not catastrophic.⁶⁷ Fuller et al⁴⁷ conducted the most comprehensive prospective cohort study on the nature of all spinal injuries with acute presentation. The incidence of spine injuries during matches was approximately 11/1000 player match hours and 0.4/1000 player training hours. The nature of spinal injuries varied between matches and training, with players more likely to sustain a cervical spine injury during matches but a lumbar spine injury during training (primarily due to weight training or running). During matches, the tackle was implicated in 37% of spinal injuries, compared with 19% in the scrum and 17% for the ruck/maul. Focussing specifically on cervical spine injuries, the tackle was implicated in 52% of injuries compared with only 12% in the scrum, highlighting the tackle as the major source of overall cervical spine injury. Player characteristics (age and anthropometrics) were not found to influence injury risk but forwards were twice as likely to sustain a spinal injury as backs. It should be noted that the scrum was a likely source of spinal injury for front row forwards, with 58% of spinal injuries resulting from the scrum and only 13% from the tackle. Thirty-three of the 35 injuries during scrummaging were sustained by front row players and only 3 of these injuries were attributed to scrum collapse.

A prospective cohort study of neck injuries (not all spine injuries) in two Australian amateur-level rugby clubs was carried out over two seasons (2006 and 2007).⁶⁸ Neck injury incidence was 6/1000 player match hours and 0.7/1000 player training hours. Forwards suffered 79% of all neck injuries, with the front row particularly susceptible, sustaining 38% of all neck injuries for only 20% of the overall player numbers and sustaining the majority of severe injuries (>3 weeks absence). Overall, the tackle was the phase of play producing most neck injuries (42%), followed by the ruck/maul (30%) and then the scrum (25%). The most common injuries were cervical facet injury (42%), followed by brachial/plexus cervical nerve root injury (stingers / burners).

353 **Chronic Degeneration Spinal Injuries**

354 Acute injuries are the most evident and quantifiable type of injury in sports/rugby injury
355 surveillance. However, repeated exposure to mechanical stresses on musculo-skeletal
356 structures may also induce sub-critical damage, and therefore the potential for long-term
357 damage to the spine due to rugby participation is an important consideration. The effects of
358 repeated exposure to scrummaging and the associated loading of the spine are very difficult
359 to detect in the short term, but likely contribute to chronic conditions. For example, front row
360 forwards are particularly prone to premature degeneration of the cervical spine, ^{17-19,69,70}
361 which may result in osteoarthritis and functional impairment, ¹⁹ with the repeated
362 microtrauma experienced during scrummaging a likely contributory cause. ¹⁷

363 Unfortunately, longitudinal data of players following retirement from the game is not currently
364 available. It has been suggested that chronic degenerative abnormalities due to repeated
365 subfailure injuries from the repeated trauma of collisions in scrummaging and tackling may
366 be underestimated and may be frequent, particularly for front row forwards, representing an
367 under-acknowledged injury issue for rugby. ⁷¹ Quinn & Winkelstein ⁷² and Panjabi ⁷³ have
368 shown that sub-catastrophic injuries to soft tissues of the spine may happen well before
369 failure limit and result in chronic pain. Repetitive (micro) traumas may generate a detrimental
370 loop in which subfailure stresses cause degeneration of intervertebral ligaments and
371 receptors, which leads to altered functioning and feedback to corrupt muscle performance.
372 This in turn produces distorted stresses in ligaments and applies abnormal loads on the
373 facet joints, accelerating degeneration and causing pain. Cervical spine degeneration may
374 be a risk factor for traumatic spinal cord injury, ¹⁹ but there is no definitive data to support
375 this presently. However, cervical spine degeneration is likely to impact upon the wellbeing of
376 players after the end of their careers.

377 Scher ⁷⁰ produced a case series comparison demonstrating greater cervical spondylosis
378 (premature degeneration) in 150 asymptomatic club rugby players compared with 150 age-
379 matched controls from the general population . Degeneration was particularly marked in front

row forwards and in the 30-35 year age group compared with 20-25 and 25-30 year groups.

¹⁹ Using radiographic evidence of cervical spine degeneration combined with clinical symptoms in a small group of professional rugby players, Hogan et al ⁶⁹ found that experienced front row rugby players (average of 23 years of playing experience) exhibited more visual evidence of general cervical spine degeneration, but that these were not necessarily accompanied by clinical symptoms or disruption to activities of daily living over and above age-matched controls. In considering these findings it is important to note that clinical status and influence on daily living activities were obtained via questionnaire and may be subject to bias due to the perceptions of what constitutes pain or symptoms.

Other studies have demonstrated narrowing of the cervical spinal canal in rugby players compared with control (non-collision) athletes, which worsens with age. ¹⁷ All asymptomatic French professional front row rugby players were assessed using static MRI imaging of the cervical spine region in seasons 2002/03 and 2003/04 and both static and dynamic MRI in seasons 2004/05 and 2005/06. ¹⁸ There was no clear difference in the medulla to canal ratio between younger and older front row players, but older players (>21 years) had a 3-fold increase in abnormalities, mainly relating to degenerative lesions. Approximately half of the sample (56 out of 127 players) presented with an anatomical abnormality and players who exhibited an abnormal medulla-to-canal ratio were also much more likely (3-fold) to exhibit anatomical abnormalities.

In a series of studies focussing on cervical spine function in rugby players, Lark and McCarthy have demonstrated that rugby forwards have impaired cervical function. This impaired function includes: reduced cervical mobility but only some reduced proprioceptive capacity (in extension) compared with rugby backs and active controls; ²⁰ reduced active cervical range of motion after a single game ^{74,75} and over the course of a season; ⁷⁵ and an inability for neck range of motion to substantially recover during an off-season despite active rehabilitation being undertaken during this period. ⁷⁶ These findings are only partially supported by a similar study ²¹ which assessed proprioceptive (head repositioning) function

in younger rugby players and found evidence of reduced repositioning ability in rugby players compared with controls, but no difference between forwards and backs. The latter point was taken to suggest that the tackle might be responsible for proprioceptive deficit rather than the scrum since backs performed similarly in the test to forwards. Imoo et al⁷⁷ further found that rugby players with previous cervical injuries had impaired static standing balance when compared with rugby players without prior cervical spine injuries.

Mechanisms of spinal injury relating to rugby scrummaging

A number of injury mechanisms which may contribute to cervical spine injury in scrummaging have been suggested. The commonly accepted notion which has guided research and opinion is that acute injuries during scrummaging normally occur through 'hyperflexion' mechanisms.⁷⁸ Increasingly, this assertion is being challenged by research suggesting that a 'buckling' mechanism is more likely.⁷⁹

'Hyperflexion' mechanism for acute spinal injury during scrummaging

Scher⁵⁸ stated that the most common mistake was for players to engage the scrum with slight flexion of the neck. This results in an elimination of the normal cervical lordosis⁸⁰ so that during a mistimed or misdirected scrum engagement or a collapsed scrum the load is applied to the flexed cervical spine rather than across the shoulders. Under this paradigm, the most common mechanism of cervical spine injury during scrummaging has been identified as hyperflexion, with or without rotation leading to anterior dislocations and unilateral or bilateral locking of facet joints.⁷⁸ McIntosh⁸¹ found the typical pattern of loading relating to neck injury was axial loading accompanied by a bending moment, a loading type that may occur during scrum engagement. The orientation of the applied load, the presence of constrained motion and the amount of energy absorbed have been found to determine the failure mode of the cervical spine, these factors all being relevant to the scrum situation. Work from Milburn¹⁴ with forward packs scrummaging against an instrumented machine has confirmed that the forces measured on engagement could be sufficient to destabilise the

spine, and more recent research has demonstrated that contemporary scrummaging produces even greater forces.^{15,25,82} Milburn⁸³ also highlighted that the bound rugby scrum places the cervical spine at risk of injury. He identified that “charging in” or misalignment of the head during engagement may result in injury, either via hyperextension (popping out) or more commonly from compression and hyperflexion of the cervical spine.

‘Buckling’ Mechanism for acute / chronic spinal injury during scrummaging

Winkelstein et al⁸⁴ suggested that an injury classification based on exceeding the range of motion e.g. hyperflexion or hyperextension is not always applicable because injury often occurs only a few milliseconds (2-20 ms) after impact when the known limits of movement of the cervical spine are still far from being reached. The contention is that a hyperflexion mechanism frequently does not explain the type of injury occurring in experimentally-induced situations, for instance there may be compression-flexion type injuries without head flexion.⁸⁵ Several authors have described the concept of a “buckling” mechanism, first introduced by Torg et al.⁸⁶ Buckling describes the mechanical instability that occurs when a structure is deformed primarily in compression, leading to changes in its deformation to a pattern of bending in compression, like compressing a long flexible ruler. This type of deformation and injury pattern has been reproduced in a number of experimental models⁸⁵ and is said to reproduce the types of injury seen in cervical spine injuries, with concurrent regions of compression alone, compression with flexion and compression with extension.

A review⁸⁵ of the biomechanics of acute cervical spine injury concluded that these compression types of injury can occur at relatively low velocities (3.1 m/s) and with relatively low loads or low percentages of total body weight (e.g. 16 kg) involved or acting on the spine.^{85,87} The risk of injury depends on a number of factors, including constraint of head-neck complex motions which would normally allow escape from the torso, and the orientation of the impact surface. The very low frequency of cervical spinal injury following head impact

has been explained by the remarkable flexibility of the neck.⁸⁸ Constraints applied to cervical motion such as “pocketing” in of the head (restricted motion of the head) are therefore thought to increase the risk of injury by increasing stiffness of the system, and preventing escape.^{85,89} With respect to impact injuries Nightingale et al⁸⁵ also showed that additional constraint of the head by the impact surface, i.e. “pocketing”, may increase the risk of injury but is not required for the injury to happen. They showed that the point of impact and the characteristics of the impacting interface has an effect on injury risk and may explain why apparently similar impacts can have dramatically different consequences. Impacts perpendicular to the cervical spine placed it at increased risk for injury compared to those where the spine's orientation was not perpendicular to the impact surface. In a neutral position, the cervical spine has a flexion lordosis of approximately 25 degrees from horizontal at T1 and it has been shown that impacts to the vertex of the head and up to 15 degrees anterior to that point have a higher frequency and severity of cervical spinal injuries than impacts anterior to this or to the posterior portion of the head.⁸⁷ This work informed the "heads-up" campaign in American football and has been attributed with reducing cervical injuries.⁸⁰ The potential role of the neck muscles in providing some protection from injury may be limited in this situation because load is mainly axial in compression and there are no muscles that resist this movement.

Despite disagreement regarding mechanism, there is general consensus that situations should be avoided where 1) spinal elements are subjected to simultaneous compression and bending loads, and 2) sudden loads are applied, since it reduces the influence of the visco-elastic elements to dampen the forces and doesn't provide time for active muscular responses.

‘Hyperflexion’ or ‘Buckling’ Mechanism for acute / chronic spinal injury during scrummaging

Kuster and colleagues ⁷⁹ conducted a systematic review of studies which considered rugby union-related cervical spine injury mechanisms and concluded that it was unlikely that the traditionally quoted hyperflexion mechanism was the true mechanism for acute injuries involving spinal cord impairment. Their interpretation was that the weight of evidence suggests the primary mechanism for the commonly observed bilateral facet joint dislocation (normally C5-C7) injury to be buckling. In opposition, Dennison et al, ⁹⁰ stated that it is too early to conclude that buckling is the predominant mechanisms of injury within the rugby union context. This opposition was partly based on the limitations associated with the *ex vivo* cadaveric testing upon which some of Kuster’s evidence was based and the fact that the same injuries produced via buckling mechanisms in cadavers have not been recreated *in vivo*, possibly due to active involvement of the musculature in protecting from injury. Therefore, there is consensus that the C4-C6 region is the most common area of injury, but the precise mechanisms for acute spinal cord injuries during scrummaging are still not clear.

Timing of acute spinal injury during scrummaging

Earlier studies which considered at which time point in the scrum injuries were sustained tended to conclude that cervical spine injuries were a result of scrum collapse. For instance, Scher ⁹¹, reported that 16 out of 40 scrum-related cervical spine injuries studied were sustained by front row forwards and reported to be due to scrum collapse. Similarly, Silver ^{63,65} reported that the vast majority of scrum-related injuries were due to collapse as opposed to engagement, and in Australian rugby between 1960 and 1996, seven scrum injuries were attributed to collapse with four attributed to engagement. ⁵⁰ Contrary to this, Wetzler ⁶⁴ analysed injury data from 1970-1996 and found a statistical difference ($P<0.002$) to demonstrate that more scrum-related cervical spine injuries occurred during engagement rather than collapse. When Quarrie et al ¹⁶ reviewed the available published data (in 2002) of

170 spinal injuries that occurred during scrummaging, an average of 47% (range 8-65%) occurred during the engagement phase, with 46% (range 29-75%) attributed to collapse. Similarly, Brown et al ⁴⁶ assimilated injury data in South Africa from 2008-2011 and reported that 56% of the scrum injuries were considered due to scrum engagement, with 39% due to scrum collapse. The differing findings across studies may reflect a changing profile of scrum-related injuries from a historical tendency for injuries to be due to scrum collapse to an increasing proportion of injuries to occur during engagement. This transition may be a reflection on the more impulsive (dynamic) nature of scrum engagement used in contemporary rugby union, which first appeared in the late 1990s.

Biomechanics of rugby scrummaging

The biomechanics of rugby scrummaging has been investigated for injury reduction/prevention (e.g. ¹⁴) and performance profiling (e.g. ²³) purposes. Most studies have employed an experimental model of one forward pack scrummaging against an instrumented scrum machine, allowing good experimental control and better repeatability than live scrummaging, but not replicating the conditions of live scrummaging. Generally, the literature indicates that rugby scrummaging involves an initial impact-like engagement phase followed by a more steady-state sustained push phase. The majority of force is produced in a forward (compression) direction but the magnitude of shear forces in the vertical direction can be considerable and lateral forces also exist. The forces produced in scrummaging have been sporadically measured over the last 25 years with a general trend for more recent studies to demonstrate greater magnitudes of force production (Table 2).

Application of forward forces

Milburn ¹⁴ investigated the forces applied by forward packs scrummaging on a rigid instrumented scrum machine. The magnitude of summed forward forces during the

engagement phase ranged from 4430 N (high school) to 7982 N (international). The observed impulsive forces were due to the large masses and 'high' speeds involved, and therefore assumed to be due mainly to the momentum generated by speed of engagement rather than active muscle action on impact. Considering primarily the forces produced by individuals and entire forward packs during sustained scrummaging, Quarrie & Wilson ²³ reported the mean sustained force from seven Community/Elite packs to be 7170 N. The sum of the force produced by each individual in each forward pack was also measured during individual scrummaging and the force produced by teams was on average 65% of the sum of these individual forces. Those packs that generated the largest scrum force were those that managed to use individual scrummaging forces to the greatest extent, thus emphasising the requirement for teams to develop technique and coordination as a unit in order to maximise pushing force.

Preatoni et al ¹⁵ described the characteristic compression force curve (Figure 2) from scrum machine trials on a range of playing levels, with the short-duration impact peak, a drop in force to a minimum level, before a gradual rise to a relatively steady-state sustained push force. The mean peak compression forces during engagement ranged from 8700 N (Women) to 16500 N (Elite and International), whilst average sustained forces ranged from 4800 N (Women) to 8300 N (International). When forces were normalised by summed body weight there was no differences in peak engagement force between Community, Academy, Women and School playing levels, but International and Elite levels still produced more force, indicative of an overall more dynamic style of scrummaging in these playing levels even accounting for body mass.

Du Toit et al ⁹² employed a novel measurement approach for measuring forces during live scrummaging via the use of pressure transducers attached to the shoulders of each player. This study recorded a maximum engagement force of approximately 10 kN (10,000 N) across an under 19 front row when they engaged with an opposition pack (so two packs generating engagement speed rather than one pack against a static scrum machine). On

average, the forces applied by the front rows during sustained scrummaging were significantly lower in magnitude than during engagement ($P<0.01$), although in one-off trials these magnitudes were very similar. Similar to Milburn¹⁴, this study found engagement forces to be positively related to the combined mass of the opposing packs, although this correlation was not present during sustained scrummaging, therefore suggesting that technique plays more of a role during the sustained phase. Cazzola et al²⁶ provided a recent measurement of live scrummaging mechanics, recording mean peak engagement forces of 9.8 kN in a sample of professional senior players.

**** Figure 2 here ****

Application of vertical and lateral shear forces

Given that the direction of movement towards the engagement is primarily horizontal and after this the primary aim of scrummaging is to push the opposing pack backward, it would be expected that the compression component of force would be the largest and the magnitude of the shear forces relatively much smaller. Milburn¹⁴ reported downward forces (~1000 N, up to 20% of the compression force value magnitude) during the engagement phase in all playing levels except for International level. It was suggested that the destabilising moment caused by the downward force would be resisted by leg extension actions of the front row players but that the presence of the downward forces would heighten the risk of collapse. Retiere⁸² reported a similar magnitude (~1500 N, approximately 12% of the peak compression force magnitude) of downward forces in the engagement phase for the French U19 team. Preatoni et al¹⁵ however, reported downward forces of greater magnitude during the engagement phase, ranging from -2000 N for School level to -3900 N for International packs (24% of the peak compression force magnitude), with a gradual transition to a slight upward force during the sustained phase. It seems plausible that the magnitude of downward force observed from machine scrummaging is in part a function of

the design of the scrum machine and the amount of downward pressure players feel confident exerting onto it.

The presence of lateral shear forces during both the engagement and sustained phases of scrummaging were highlighted by Milburn as being inefficient and, over the long term, a likely cause of premature degeneration of the cervical spine.⁷⁰ The proposed mechanism is that shear forces introduce a moment of force which is not present during pure compression and which induces undesirable rotation and/or bending of the spine. Preatoni et al¹⁵ found the patterns of lateral forces during engagement to be lower in magnitude than compression and vertical forces (approximately 10% of compression force magnitude) and inconsistent in direction.

Given the values reported in the different studies it appears that the forces involved in rugby scrummaging have increased considerably in the last twenty years, particularly during the engagement phase. These changes may be due to a combination of increased player size and a more dynamic engagement action, although differences in experimental instrumentation (e.g. more rigid scrum machine structures used in older studies) should not be ruled out as a contributing factor. In support of the suggestion that the engagement process has become more dynamic, the speed of engagement of International-level packs in Preatoni et al's study¹⁵ (~3.0 m/s, in 2013) was considerably greater than the engagement speed observed in Milburn's 1990 study¹⁴ (~2.0 m/s, in 1990).

611 Table 2. Forces generated during rugby scrummaging

Study	Playing Level	Engagement			Sustained			Study Details
		Peak Forward / Compression (N)	Peak Vertical (N)	Peak Lateral (N)	Average Forward / Compression (N)	Average Vertical (N)	Average Lateral (N)	
Milburn ¹⁴	School	4430	-940	-150	3370	190	-3040	Scrum Machine; Rigid frame; 500 Hz sampling; 1 team per level
	University	6540	-160	-730	4610	610	-1510	
	Community	5630	-868	-2413	4300	-151	-3093	
	International	7982	2268	-85	5761	1305	-340	
Rodano & Tosoni ²⁴	International U19	~11400	~4400	~400				Rigid frame; 500 Hz sampling; Single players and simulated pack reconstruction
Quarrie & Wilson ²³	Community	11000			7170			Scrum machine; 20 Hz sampling; Forces represent absolute (modulus) of force
Du Toit ⁹²	School	7526			6145			Live scrum; Pressure transducers; Force derived from summed pressures Mean of 13 teams
Retiere ⁸²	International U19	~ 12000	~ -1500		~ 7000	~ -200		Scrum machine; 500 Hz sampling; Damping in machine pads; 1 team
Preatoni ¹⁵	School	9100	-2000	1100	4880	100	110	Scrum machine; 500 Hz sampling; Damping in machine pads; 4-6 teams

University	11700	-2900	1300	5940	96	130	per level; Lateral force during engagement is range of lateral force not peak.
Women	8700	-2400	1000	4790	7	-90	
Community	12000	-2300	1400	5780	-28	110	
Elite Club	16500	-3900	1900	8300	720	620	
International	16500	-3600	1900	8300	1084	600	

612

613

614 ***Modifying the engagement process***

615 One potential route to modifying the forces generated in the rugby scrum, particularly during
616 the initial engagement phase, is to modify the engagement process through changes to
617 player actions or referee instructions. Previously this has been attempted by adjusting how
618 many players are involved in the engagement. Milburn & O'Shea ⁹³ investigated a sequential
619 scrum formation whereby, each row of the scrum formed once the previous row were in
620 position (i.e., 3+2+3). Sequential engagement significantly reduced the total engagement
621 force (4833 N) experienced across the front row players compared with the standard scrum
622 (5882 N, $P<0.05$), primarily due to a reduction in force through the loose head prop.
623 However, sequential formation also reduced the stability of the scrum by increasing its
624 duration and increasing the variability of vertical and lateral forces acting on the front row
625 players as second rows and back row players were added asynchronously. Similarly, Du Toit
626 et al ⁹² found that the peak engagement force of all players in the scrum was significantly
627 greater during a full scrum engagement (7526 N) as opposed to a sequential scrum
628 engagement (4596 N, $P<0.01$) in under 19 schoolboy teams, with no differences in the
629 forces achieved during sustained scrummaging. Retiere ⁸² reported downward forces for the
630 French U19 team in the region of -1000 N for a normal engagement and -600 N for a
631 sequential 5+3 engagement, where the front and second row only were involved in the initial
632 engagement and the back row were added subsequently.

633 In machine-scrummaging trials, Preatoni et al ²⁵ demonstrated that a 5+3 sequential
634 engagement significantly ($P<0.05$) reduced peak compression (12-20%) and downward (5-
635 32%) forces during the engagement phase for all playing levels but did not alter a 'Hazard
636 Index' combining measures of force and head-neck alignment. However, a 'fold-in'
637 engagement where all 8 forwards were involved but were instructed to de-emphasise the
638 engagement created larger reductions in peak compression forces (45-54%) and downward
639 forces (21-40%) as well as significantly reducing the Hazard Index measure ($P<0.05$). This

fold-in procedure also allowed forward packs to maintain forward force generation during the sustained push phase.

Measuring player loading variables in the more realistic setting of contested live scrummaging, Cazzola et al ²⁶ performed an initial study in a group of elite professional teams, demonstrating an approximate reduction of peak forces across the front row during engagement of approximately 25% when using a pre-bind engagement process (~6300 N) compared with the 2012-13 full scrum engagement process (~8800 N). This pre-bind engagement process did not impair force generation in the sustained phase of scrummaging and also did not negatively influence scrum stability measures.

In summary, a number of studies have shown that a sequential engagement process for the scrum, by progressively adding players following the initial engagement of the two forward packs in some way, reduces peak forces experienced by front row players but upsets the stability of the scrum in terms of creation of shear forces, spinal misalignments, or overall duration of the scrum. Therefore, the principle of sequential scrum engagement has not been recommended by any of the published studies. On the other hand, engagement processes which involve the full scrum configuration (all 16 players) in the initial engagement phase but which de-emphasises the momentum generated during this phase appear to produce more encouraging results in terms of force reduction alongside maintenance of scrum stability.

CONCLUSION

This review has highlighted that, scrummaging accounts for up to, but probably no more than, 10% of all rugby-related injuries. Most of these reported injuries are of moderate severity and the incidence of catastrophic injuries from scrummaging is very low. Approximately 40% of all rugby-related spinal cord injuries can be attributed to the scrum. Conclusive statements regarding the true level and trends of catastrophic injuries in rugby union have been hampered by a lack of consistency coherence in medical record keeping

667 and poor estimates of the size of the rugby-playing population. In recent years the
668 International Rugby Board has constituted a centralised database intended to capture all
669 catastrophic injuries occurring world-wide and so a clearer picture should become apparent.

670 There is also emerging evidence regarding the issue of chronic degeneration in rugby
671 players, with the suggestion that scrummaging may play a role in the deleterious anatomical
672 and functional effects displayed by rugby forwards. Again, a lack of longitudinal clinical
673 datasets on cohorts of rugby players and matched controls makes definitive statements
674 around the influence of rugby, and scrummaging in particular, on degeneration of the spine
675 difficult to make and this is a key area for future research.

676 During the engagement phase, the forces generated at the interface between the two front
677 rows during scrummaging are considerable and include forces in multiple directions, mainly
678 forward but also downward. The forces acting during engagement can be modified but
679 negative consequences in terms of stability have been reported when sequential scrum
680 engagement processes have been attempted; limitations apparently not observed when fold-
681 in/pre-bind engagement processes are employed.

682 The relatively “controlled” environment of the scrum is a phase of play in which it should be
683 possible to intervene to reduce injury occurrence, either through modifications to player
684 technique, coaching practices or laws.¹⁰ The scrum therefore remains high priority for
685 research with a long term goal of injury reduction.

686

687 **COMPETING INTERESTS**

688 None of the authors has competing financial, professional or personal interests that might
689 have influenced the performance or presentation of the work described in this manuscript.

690

691 **FUNDING**

692 This project is funded by the International Rugby Board (IRB).

693

694

REFERENCE LIST

1. Roberts SP, Trewartha G, Higgitt RJ, et al. The physical demands of elite English rugby union. *J Sports Sci* 2008;26(8):825-33.
2. Brooks JH, Fuller CW, Kemp SP, et al. Epidemiology of injuries in English professional rugby union: part 1 match injuries. *Br J Sports Med* 2005;39(10):757-66.
3. Fuller C, Laborde F, Leather R, et al. International Rugby Board Rugby World Cup 2007 injury surveillance study. *Br J Sports Med* 2008;42:452 - 9.
4. Fuller CW, Raftery M, Readhead C, et al. Impact of the International Rugby Board's experimental law variations on the incidence and nature of match injuries in southern hemisphere professional rugby union. *South Afr Med J* 2009;99(4):232-7.
5. Williams S, Trewartha G, Kemp S, et al. A Meta-Analysis of Injuries in Senior Men's Professional Rugby Union. *Sports Med Online First*: doi: 10.1007/s40279-013-0078-1.
6. Gissane C, Jennings D, Kerr K, et al. A pooled data analysis of injury incidence in rugby league football. *Sports Med*. 2002;32(3):211-16.
7. Kaplan LD, Flanigan DC, Norwig J, et al. Prevalence and variance of shoulder injuries in elite collegiate football players. *Am J Sports Med*. 2005;33(8):1142-46.
8. Lorentzon R, Wedren H, Pietila T, et al. Injuries in International ice hockey - a prospective, comparative-study of injury incidence and injury types in International and Swedish elite ice hockey. *Am J Sports Med* 1988;16(4):389-91.

- 719 9. Orchard J, Seward H. Epidemiology of injuries in the Australian Football League,
720 seasons 1997-2000. *Br J Sports Med* 2002;36(1):39-44.
- 721 10. Raftery M. The past, the present and the future. International Rugby Board
722 Medical Conference; 21/11/2011; London, UK.
- 723 11. Fuller CW. Catastrophic injury in rugby union is the level of risk acceptable?
724 *Sports Med* 2008;38(12):975-86.
- 725 12. Bourke JB. Rugby union should ban contested scrums. *BMJ*
726 2006;332(7552):1281-81.
- 727 13. IRB. <http://www.irblaws.com/index.php?law=20>. International Rugby Board, 2013
728 (date accessed May 15th 2013).
- 729 14. Milburn PD. The kinetics of rugby union scrummaging. *J Sports Sci* 1990;8(1):47-
730 60.
- 731 15. Preatoni E, Stokes KA, England ME, et al. The influence of playing level on the
732 biomechanical demands experienced by rugby union forwards during machine
733 scrummaging. *Scand J Med Sci Sports* 2013;23:e178-e184.
- 734 16. Quarrie KL, Cantu RC, Chalmers DJ. Rugby Union injuries to the cervical spine
735 and spinal cord. *Sports Med* 2002;32(10):633-53.
- 736 17. Berge J, Marque B, Vital JM, et al. Age-related changes in the cervical spines of
737 front-line rugby players. *Am J Sports Med* 1999;27(4):422-9.
- 738 18. Castinel BH, Adam P, Milburn PD, et al. Epidemiology of cervical spine
739 abnormalities in asymptomatic adult professional rugby union players using
740 static and dynamic MRI protocols: 2002 to 2006. *Br J Sports Med*
741 2010;44(3):194-99.
- 742 19. Scher AT. Premature onset of degenerative disease of the cervical spine in
743 rugby players. *South Afr Med J* 1990;77(11):557-8.

- 744 20. Lark SD, McCarthy PW. Cervical range of motion and proprioception in rugby
745 players versus non-rugby players. *J Sports Sci* 2007;25(8):887-94.
- 746 21. Pinsault N, Anxionnaz M, Vuillerme N. Cervical joint position sense in rugby
747 players versus non-rugby players. *Phys Ther Sport* 2010;11(2):66-70.
- 748 22. Du Toit DE, Venter DJL, Buys FJ, et al. Kinetics of rugby union scrumming in
749 under-19 schoolboy rugby forwards. *South African Journal for Research in*
750 *Sport, Physical Education and Recreation* 2004;26(2):33-50.
- 751 23. Quarrie KL, Wilson BD. Force production in the rugby union scrum. *J Sports Sci*
752 2000;18(4):237-46.
- 753 24. Rodano R, Tosoni A. *La mischia nel rugby*. Milano: Edi Ermes s.r.l, 1992.
- 754 25. Preatoni E, Stokes KA, England M, et al. Engagement techniques and playing
755 level impact the biomechanical demands on rugby forwards during machine-
756 based scrummaging. *Br J Sports Med* 2013;under review.
- 757 26. Cazzola D, Preatoni E, Stokes KA, et al. A modified pre-bind engagement
758 process reduces biomechanical loading on front row players during
759 scrummaging: a pilot study of 11 elite teams. *Br J Sports Med* 2013;under
760 review.
- 761 27. Brooks JH, Kemp SP. Recent trends in rugby union injuries. *Clin Sports Med*
762 2008;27(1):51-73, vii-viii.
- 763 28. Kerr H, Curtis C, Micheli L, et al. Collegiate rugby union injury patterns in New
764 England: a prospective cohort study. *Br J Sports Med* 2008;42:595-603.
- 765 29. Roberts SP, Trewartha G, England M, et al. Epidemiology of time-loss injuries in
766 English community level rugby union. *BMJ Open* 2013;accepted for
767 publication.

- 768 30. Schneiders AG, Takemura M, Wassinger CA. A prospective epidemiological
769 study of injuries to New Zealand premier club rugby union players. *Phys Ther*
770 *Sport* 2009;10(3):85-90.
- 771 31. Haseler CM, Carmont MR, England M. The epidemiology of injuries in English
772 youth community rugby union. *Br J Sports Med* 2010;44(15):1093-99.
- 773 32. Nicol A, Pollock A, Kirkwood G, et al. Rugby union injuries in Scottish schools. *J*
774 *Public Health* 2011;33(2):256-61.
- 775 33. Palmer-Green DS, Stokes KA, Fuller CW, et al. Match injuries in English youth
776 academy and schools rugby union: an epidemiological study. *Am J Sports*
777 *Med* 2013;41(4):749-55.
- 778 34. Roi GS, Bonanno D, Pacini D, et al. Youth rugby union injury rates and
779 suggestions for injury prevention. *Medicina Dello Sport* 2010;63(4):567-77.
- 780 35. Schick DM, Molloy MG, Wiley JP. Injuries during the 2006 Women's Rugby
781 World Cup. *Br J Sports Med* 2008;42(6):447-51.
- 782 36. Taylor AE, Fuller CW, Molloy MG. Injury surveillance during the 2010 IRB
783 Women's Rugby World Cup. *Br J Sports Med* 2011;45(15):1243-45.
- 784 37. Targett SG. Injuries in professional Rugby Union. *Clin J Sport Med*
785 1998;8(4):280-5.
- 786 38. Taylor AE, Kempt SPT, Trewartha G, et al. Scrum injury risk in professional
787 rugby union. *Br J Sports Med* 2013;under review.
- 788 39. Palmer-Green D. Injury epidemiology and injury prevention in English youth
789 rugby union. Unpublished PhD Thesis, University of Bath, 2010.
- 790 40. Fuller CW, Brooks JHM, Cancea RJ, et al. Contact events in rugby union and
791 their propensity to cause injury. *Br J Sports Med* 2007;41(12):862-67.

- 792 41. Roberts SP, Trewartha G, England M, et al. Propensity of contact events to
793 cause injury in community rugby union: a focus on collapsed scrums and
794 collision tackles. *Br J Sports Med* 2013;under review.
- 795 42. The incidence, severity and nature of scrummaging injuries in professional rugby
796 union. 1st World Congress on Sports Injury Prevention; 2005.
- 797 43. Brooks JHM, Kemp SPT. Injury-prevention priorities according to playing position
798 in professional rugby union players. *Br J Sports Med* 2011;45(10):765-75.
- 799 44. Torg JS, Vegso JJ, Sennett B. The National Football Head and Neck Injury
800 Registry: 14-year report on cervical quadriplegia (1971-1984). *Clin Sports*
801 *Med* 1987;6(1):61-72.
- 802 45. IRB. <http://www.irbplayerwelfare.com/?documentid=11&language=en>,
803 International Rugby Board, 2013.
- 804 46. Brown JC, Lambert MI, Verhagen E, et al. The incidence of rugby-related
805 catastrophic injuries (including cardiac events) in South Africa from 2008 to
806 2011: a cohort study. *BMJ Open* 2013;3(2): doi: pii: e002475.
807 10.1136/bmjopen-2012-002475..
- 808 47. Fuller CW, Brooks JH, Kemp SP. Spinal injuries in professional rugby union: a
809 prospective cohort study. *Clin J Sport Med* 2007;17(1):10-6.
- 810 48. Maharaj JC, Cameron ID. Increase in spinal injury among rugby union players in
811 Fiji. *Med J Aust* 1998;168(8):418.
- 812 49. Rotem TR, Lawson JS, Wilson SF, et al. Severe cervical spinal cord injuries
813 related to rugby union and league football in New South Wales, 1984-1996. *M*
814 *J Aust* 1998;168(8):379-81.
- 815 50. Taylor TKF, Rutkowski SB, Jones RF, et al. Spinal cord injuries in Australian
816 footballers. *ANZ Journal of Surgery* 2003;73(7):493-99.

- 817 51. Van Mechelen W. The severity of sports injuries. *Sports Med*. 1997;24:176 - 80.
- 818 52. Bohu Y, Julia M, Bagate C, et al. Declining Incidence of Catastrophic Cervical
819 Spine Injuries in French Rugby 1996-2006. *Am J Sports Med* 2009;37(2):319-
820 23.
- 821 53. Hermanus FJ, Draper CE, Noakes TD. Spinal cord injuries in South African
822 Rugby Union (1980-2007). *South Afr Med J* 2010;100(4):230-34.
- 823 54. Carmody DJ, Taylor TKF, Parker DA, et al. Spinal cord injuries in Australian
824 footballers 1997-2002. *Med J Aust* 2005;182(11):561-64.
- 825 55. MacLean JG, Hutchison JD. Serious neck injuries in U19 rugby union players: an
826 audit of admissions to spinal injury units in Great Britain and Ireland. *Br J*
827 *Sports Med* 2012; 46: 591-94.
- 828 56. Berry JG, Harrison JE, Yeo JD, et al. Cervical spinal cord injury in rugby union
829 and rugby league: are incidence rates declining in NSW? *ANZ J Public Health*
830 2006;30(3):268-74.
- 831 57. Noakes TD, Jakoet I, Baalbergen E. An apparent reduction in the incidence and
832 severity of spinal cord injuries in schoolboy rugby players in the Western
833 Cape since 1990. *South Afr Med J* 1999;89(5):540-45.
- 834 58. Scher AT. Rugby injuries to the cervical spine and spinal cord: A 10-year review.
835 *Clin Sports Med* 1998;17(1):195-206.
- 836 59. Bottini E, Poggi EJT, Luzuriaga F, et al. Incidence and nature of the most
837 common rugby injuries sustained in Argentina (1991-1997). *Br J Sports Med*
838 2000;34(2):94-97.
- 839 60. Secin FP, Poggi EJT, Luzuriaga F, et al. Disabling injuries of the cervical spine in
840 Argentine rugby over the last 20 years. *Br J Sports Med* 1999;33(1):33-36.

- 841 61. Kew T, Noakes TD, Kettles AN, et al. A retrospective study of spinal cord injuries
842 in Cape Province rugby players, 1963-1989. Incidence, mechanisms and
843 prevention. *South Afr Med J* 1991;80(3):127-33.
- 844 62. Silver JR. The impact of the 21st century on rugby injuries. *Spinal Cord*
845 2002;40(11):552-59.
- 846 63. Silver JR, Gill S. Injuries of the spine sustained during rugby. *Sports Med*.
847 1988;5(5):328-34.
- 848 64. Wetzler MJ, Akpata T, Laughlin W, et al. Occurrence of cervical spine injuries
849 during the rugby scrum. *Am J Sports Med* 1998;26(2):177-80.
- 850 65. Silver JR. Injuries of the spine sustained in rugby. *BMJ* 1984;288(6410):37-43.
- 851 66. Taylor TKF, Coolican MRJ. Spinal-cord injuries in Australian footballers, 1960-
852 1985. *Med J Aust* 1987;147(3):112-118.
- 853 67. McIntosh AS, McCrory P, Finch CF, et al. Head, face and neck injury in youth
854 rugby: incidence and risk factors. *Br J Sports Med* 2010;44(3):188-93.
- 855 68. Swain M, Pollard H, Bonello R. Incidence, severity, aetiology and type of neck
856 injury in men's amateur rugby union: a prospective cohort study. *Chiropractic*
857 *& Osteopathy* 2010;18(1):18.
- 858 69. Hogan BA, Hogan NA, Vos PM, et al. The cervical spine of professional front-row
859 rugby players: correlation between degenerative changes and symptoms. *Ir J*
860 *Med Sci* 2010;179(2):259-63.
- 861 70. Scher AT. Serious cervical spine injury in the older rugby player. An indication for
862 routine radiological examination. *South Afr Med J* 1983;64(4):138-40.
- 863 71. Yeo JD. Rugby and spinal injury: what can be done? *Med J Aust*
864 1998;168(8):372-73.

- 865 72. Quinn KP, Winkelstein BA. Cervical facet capsular ligament yield defines the
866 threshold for injury and persistent joint-mediated neck pain. *J Biomech*
867 2007;40(10):2299-306.
- 868 73. Panjabi M. A hypothesis of chronic back pain: ligament subfailure injuries lead to
869 muscle control dysfunction. *Eur Spine J* 2006;15(5):668-76.
- 870 74. Lark SD, McCarthy PW. The effects of a single game of rugby on active cervical
871 range of motion. *J Sports Sci* 2009;27(5):491-97.
- 872 75. Lark SD, McCarthy P. The effects of a rugby playing season on cervical range of
873 motion. *J Sports Sci* 2010;28(6):649-55.
- 874 76. Lark S, McCarthy P. Active cervical range of motion recovery following the rugby
875 off-season. *J Sports Med Phys Fitness* 2010;50(3):318-25.
- 876 77. Imoo Y, Takemura M, Takuo F, et al. Standing balance ability of Japanese
877 collegiate rugby union players with past cervical injuries. *Football Science*
878 2013;10:1-9.
- 879 78. Scher AT. Crashing the Rugby Scrum - an Avoidable Cause of Cervical Spinal-
880 Injury. *South Afr Med J* 1982;61(24):919-20.
- 881 79. Kuster D, Gibson A, Abboud R, et al. Mechanisms of cervical spine injury in
882 Rugby Union: a systematic review of the literature. *Br J Sports Med*
883 2012;46:550-54.
- 884 80. Torg JS, Vegso JJ, Oneill MJ, et al. The epidemiologic, pathological,
885 biomechanical, and cinematographic analysis of football-induced cervical-
886 spine trauma. *Am JSports Med* 1990;18(1):50-7.
- 887 81. McIntosh AS. Rugby injuries. *Med Sport Sci* 2005;49:120-39.
- 888 82. Retiere D. Expérimentations simulateur de mêlée pole France. International
889 Rugby Board Medical Conference; 04/11/2010; London, UK.

- 890 83. Milburn PD. Biomechanics of Rugby Union Scrummaging *Sports Med.*
891 1993;16(3):168-79.
- 892 84. Winkelstein B, Myers B. The biomechanics of cervical spine injury and
893 implications for injury prevention. *Med Sci Sports Exerc* 1997;29:246-55.
- 894 85. Nightingale RW, McElhaney JH, Richardson WJ, et al. Dynamic responses of the
895 head and cervical spine to axial impact loading. *J Biomech* 1996;29(3):307-
896 18.
- 897 86. Torg JS, Pavlov H, O'Neill MJ, et al. The axial load teardrop fracture. A
898 biomechanical, clinical and roentgenographic analysis. *Am J Sports Med*
899 1991;19(4):355-64.
- 900 87. Nightingale RW, Richardson WJ, Myers BS. The Effects of Padded Surfaces on
901 the Risk for Cervical Spine Injury. *Spine* 1997;22(20):2380-87.
- 902 88. Roaf R. A study of the mechanics of spinal injuries. *J Bone Joint Surg*
903 1960;42B(4):810-23.
- 904 89. Yoganandan N, Sances A, Maiman DJ, et al. Experimental spinal-injuries with
905 vertical impact. *Spine* 1986;11(9):855-60.
- 906 90. Dennison CR, Macri EM, Cripton PA. Mechanisms of cervical spine injury in
907 rugby union: is it premature to abandon hyperflexion as the main mechanism
908 underpinning injury? *Br J Sports Med* 2012;46(8):545-49.
- 909 91. Scher AT. Catastrophic rugby injuries of the spinal cord: changing patterns of
910 injury. *Br J Sports Med* 1991;25(1):57-60.
- 911 92. Du Toit DE, Olivier PE, Buys FJ. Kinetics of full scrum and staggered scrum
912 engagement in under-19 schoolboy rugby union players. *South African*
913 *Journal for Research in Sport, Physical Education and Recreation*
914 2005;27(2):15-28.

915 93. Milburn PD, O'Shea BP. The sequential scrum engagement: A biomechanical
916 analysis. *Aust J Science Med Sport* 1994;26(1):32-35.

917

918

919

FIGURE CAPTIONS

Figure 1. Key events and body postures during rugby scrum engagement. Following a referee call of “crouch” then “touch” the front rows crouch so that when they meet, each player’s head and shoulders are no lower than the hips and props touch the shoulder of the opposing prop before withdrawing their arm. The referee then calls “set” (as of August 2012), which is an indication that the front rows may come together when ready. The front rows of each team’s scrum pack engage with their heads interlocked, with contact between the front row players taking place through the backs of their necks and shoulders. As a result of this a tunnel is created into which the scrum-half throws in the ball and the forward packs compete for possession by aiming to push the opposing pack backwards.

Figure 2. Characteristic force traces typical of those obtained from studies involving one forward pack scrummaging against an instrumented scrum machine, adapted from Preatoni et al ¹⁵

Crouch Position



Touch Position




Set Position (post-engagement)



936 Figure 1

937



938 Figure 2